

PATENT SPECIFICATION

(11) 1 515 425

1 515 425

- (21) Application No. 20639/76 (22) Filed 19 May 1976
 (21) Application No. 49608/76 (22) Filed 29 Nov. 1976
 (23) Complete Specification filed 28 April 1977
 (44) Complete Specification published 21 June 1978
 (51) INT CL² A23L 3/34; A21D 2/14
 (52) Index at acceptance A2D 2L 3B2X 3B4X 3BX
 (72) Inventor DENNIS JAMES GEORGE LONG

(19)



(54) PRESERVATION PROCESS

PATENTS ACT 1949

SPECIFICATION NO 1515425

The following amendments were allowed under Section 29 on 8 December 1982:

- Page 1, lines 49 and 56, Page 5, lines 22 and 23, *after* therein *insert* an aqueous solution of
- Page 1, line 50, Page 5, lines 23 and 24, *after* defined *insert* containing the acid propionate salt
- Page 1, line 57, Page 5, line 28, *after* salt *insert* containing the acid propionate salt
- Page 1, line 82, *after* The *insert* aqueous solution of
- Page 1, line 95, *delete* preferably
- Page 2, lines 2 and 3, *delete* may be added as a solid or as *insert* is added as
- Page 5, line 28, *after* bread *insert* an aqueous solution of
- Page 5, *delete* lines 44 to 46
- Page 5, line 47, *after* wherein *insert* the aqueous solution of
- Page 5, *for* claims 7 to 9 *read* 6 to 8
- Page 5, line 57, *for* 8 *read* 7

THE PATENT OFFICE

21 March 1983

- 35 *delete* the acetates and malates of sodium. Propionates of sodium and calcium have also been claimed and described in British Patent Specification Serial No. 488,560 (Ward Baking Company). Since the principal active constituent is believed to be the acid component, the use of fully neutralised salts results in the need for adding a larger amount of the salt than should be necessary based on the acid content of the salt. It has now been found that by using the acid salts, the desired level of inhibition activity can be achieved using a relatively smaller proportion of the acid salt.
- 40
- 45 Accordingly, the present invention is a method of inhibiting the growth of harmful micro-organisms in foodstuff susceptible thereto comprising incorporating therein an acid

Bas 95178/6

The acid propionates are suitably prepared by adding a full propionate salt to propionic acid and water. For example a 70% by weight solution of sodium acid propionate may be prepared by adding sodium propionate to propionic acid in water. 85

Alternatively a solution may be prepared by the addition of solutions of sodium hydroxide or carbonate to propionic acid. For example 95 pts. of a 42% w/w aqueous solution of sodium hydroxide was added to 148 pts. of stirred, cooled propionic acid to yield 243 pts. of a 70% solution of sodium dipropionate. 90

The acid propionate salts are preferably added in an amount of between 0.01 and 0.4% by the dry weight of the foodstuff treated, but not so as to contravene any legal restrictions on 95

PATENT SPECIFICATION

(11) 1 515 425

1 515 425

- (21) Application No. 20639/76 (22) Filed 19 May 1976
 (21) Application No. 49608/76 (22) Filed 29 Nov. 1976
 (23) Complete Specification filed 28 April 1977
 (44) Complete Specification published 21 June 1978
 (51) INT CL² A23L 3/34; A21D 2/14
 (52) Index at acceptance A2D 2L 3B2X 3B4X 3BX
 (72) Inventor DENNIS JAMES GEORGE LONG



(54) PRESERVATION PROCESS

(71) We, BP CHEMICALS LIMITED, of Britannic House, Moor Lane, London, EC2Y 9BU, a British Company, do hereby declare the invention, for which we pray that a patent may be granted to us, and the method by which it is to be performed, to be particularly described in and by the following statement:—

The present invention relates to a method of inhibiting the growth of harmful micro-organisms, for instance, bacteria and fungi in foodstuffs and other organic material susceptible thereto.

Microbiological deterioration of bread and bread products is well known. The wastage of bread due to moulds and bacterial organisms has been considerable. This is basically due to the speed of production, rapid cooling and the tendency to produce softer bread which gives rise to marginally higher moisture content in the bread. Additionally, moisture retention is enhanced by the current trends in the packaging of sliced wrapped bread. This in turn renders the bread susceptible to attack by moulds and bacteria.

Various methods have hitherto been used to inhibit the growth of harmful micro-organisms. The primary consideration for the desired additive being that it attacks the micro-organism responsible without being toxic to human beings. Examples of additives used hitherto with varying degrees of success include the acetates and diacetates of sodium. Propionates of sodium and calcium have also been claimed and described in British Patent Specification Serial No. 488,560 (Ward Baking Company). Since the principal active constituent is believed to be the acid component, the use of fully neutralised salts results in the need for adding a larger amount of the salt than should be necessary based on the acid content of the salt. It has now been found that by using the acid salts, the desired level of inhibition activity can be achieved using a relatively smaller proportion of the acid salt.

Accordingly, the present invention is a method of inhibiting the growth of harmful micro-organisms in foodstuff susceptible thereto comprising incorporating therein an acid

propionate salt as hereinafter defined in an amount of between 0.01 to 1.0% by the dry weight of the foodstuff.

According to a preferred embodiment, the present invention is a method of inhibiting rope formation and mould growth in bread comprising incorporating therein an acid propionate salt in an amount of between 0.01 and 0.4% by the dry weight of the flour. The method of the present invention is particularly suitable for inhibiting rope formation and mould growth in enzyme leavened or chemically aerated bread.

Other foodstuffs that may be treated according to the present invention include products such as cakes, pastry mixes, biscuits, flour, starches, precooked foods such as sausages and meat pies and baby foods.

The "acid propionate salts" according to the present invention are the acid salts of propionic acid and at least one cation selected from ammonium, sodium, potassium, calcium and magnesium such that the ratio of acid to the cation in the salt is in the range of 2:1 to 4:1 on a chemical equivalent basis. In effect, these are mixtures of propionic acid with a neutral salt thereof e.g. sodium propionate, in at least equimolar proportions. Preferred acid salts are sodium dipropionate, ammonium dipropionate, potassium dipropionate, calcium tetrapropionate and magnesium tetrapropionate.

The acid propionates are suitably prepared by adding a full propionate salt to propionic acid and water. For example a 70% by weight solution of sodium acid propionate may be prepared by adding sodium propionate to propionic acid in water.

Alternatively a solution may be prepared by the addition of solutions of sodium hydroxide or carbonate to propionic acid. For example 95 pts. of a 42% w/w aqueous solution of sodium hydroxide was added to 148 pts. of stirred, cooled propionic acid to yield 243 pts. of a 70% solution of sodium dipropionate.

The acid propionate salts are preferably added in an amount of between 0.01 and 0.4% by the dry weight of the foodstuff treated, but not so as to contravene any legal restrictions on

1 515 425A - SEE ATTACHED SLIP

the use of preservatives in foodstuffs.

5 The acid propionate salt may be added as
a solid or as an aqueous solution to the in-
gredients before, during or after the mixing
of the ingredients depending upon the food-
stuff being treated. Furthermore, if the in-
hibitor is to be incorporated in the processing
of bread, it is preferably added to the water
used in making dough from flour rather than
10 to the solid components of the bread mix in
the bread making process.

15 The acid propionate salts are more water
soluble than the corresponding neutral pro-
pionates. This enables solutions to be prepared
and used which are sufficiently concentrated
to provide significant economic advantage over
the solid neutral propionate salt or aqueous
solutions of the neutral propionate salt. Solu-
bilities of the calcium and sodium acid and
20 neutral propionates in water are given below:

solubility at 20°C

sodium propionate	99.5 g/100 g water
sodium dipropionate	233 g/100 g water

solubility at 25°C

25 calcium propionate	39 g/100 g water
calcium tetrapropionate	48 g/100 g water

The invention is further illustrated with
reference to the following Examples.

Examples 1—3.

Using a Chorleywood Breadmaking Process 30
(C.B.P.) bread recipe, doughs were made up
containing 6, 9 and 12 oz of 70% sodium di-
propionate solution per sack of flour. In all
cases, the mould inhibitors were added 35
to the doughing water. Three dough
pieces, moulded to the 4 piece shape,
were produced from each dough and
proved for a constant time of 47 min
prior to baking. The average proof height,
oven height, oven spring and loaf volume were 40
determined. A small scale organoleptic test was
carried out using a panel of 10 persons who
were asked to rate loaves from the various
treatments for odour on a scale from not evi-
45 dent=0 to extremely evident=5. The mould-
free shelf life of the various breads was deter-
mined by cutting the loaves into five thick
slices and exposing each cut surface to the
atmosphere for 5 min to allow natural con-
50 tamination with moulds to occur prior to
wrapping. The number of mouldy cut surfaces
was determined twice daily during storage at
27°C (80°F) and an approximate mould-free
shelf life estimated. The results from this test
55 are given in Table 1.

By way of comparison with the above Ex-
amples 1—3, tests (a), (b) and (c) were
carried out using bread with no inhibitor and
with calcium propionate as inhibitor. The
60 results are shown in Table 2.

TABLE 1

Treatment	Proof Ht. (cm)	Oven Ht. (cm)	Oven Spring (cm)	Loaf vol. (ml)	Average panel rating for odour	Approx. mould- free shelf life (days at 27°C)
1. Na. acid prop. 6* oz/sack**	10.75	16.00	5.25	2870	0.5	2.75
2. Na. acid prop. 9* oz/sack**	9.5	15.75	6.25	2863	1.2	3.75
3. Na. acid prop. 12* oz/sack**	9.4	14.80	5.40	2807	2.3	4.00

* weight of a 70% aqueous solution.

TABLE 2

Treatment	Proof Ht. (cm)	Oven Ht. (cm)	Oven Spring (cm)	Loaf vol. (ml)	Average panel rating for odour	Approx. mould- free shelf life (days at 27°C)
(a) Untreated control	11.0	15.75	4.75	2909	0.1	2.25
(b) Solid Ca. prop. 9 oz/sack**	10.0	15.60	5.60	2844	0.6	3.50
(c) Solid Ca. prop. 14 oz/sack**	9.75	15.50	5.75	2812	3.1	4.75

** sack contained 280 lbs. of flour.

5 At all levels the inhibitors caused underproof in the fixed proof time employed but the oven spring was increased over that of the untreated control. Loaf volume was reduced as the amount of inhibitor present increased. The reduction in volume using 14 oz/sack of calcium propionate was similar to that when 12 oz/sack of sodium dipropionate was used.

Examples 4—9.

10 Additional experiments were carried out in a commercial bakery on three different

occasions to compare the mould-free shelf life of sliced and bagged standard bread containing 70% sodium dipropionate solution. In each experiment the sodium dipropionate solution was part of the liquor used to make up the dough. Additional yeast was added to overcome any effects on loaf volume, i.e. 8 oz/sack for 7—9 oz/sack levels of both inhibitors and 12 oz/sack for 12—14 oz/sack levels of both inhibitors.

In each experiment, 22 sliced and bagged

loaves per treatment were held in an incubator at 27°C (80°F) and examined at intervals for the numbers of loaves mouldy both on the outside and cut surfaces. Loaf volume measurements were carried out on four loaves per treatment. In the initial experiments, organoleptic tests were also carried out using a panel of 30 persons who were asked to rate the odour present in the various breads on a scale from

not evident=0 to extremely evident=5. The results from these tests are summarised in Table 3.

Again by way of comparison, Examples 4-9 were repeated but adding solid calcium propionate to the dry ingredients instead of the sodium dipropionate solution being added to the dough making liquor. The results are shown in Table 4.

TABLE 3

Treatment	Average loaf specific volume	Average panel rating for odour	No. of loaves out of 22 mouldy after					Approx. mould-free shelf life (days at 27 °C)
			3 3/4	4	4 1/4 days	5 1/4	6 1/4	
4. Na. acid prop. 9** oz/sack	3.5	1.3	0(0)	0(0)	2(1)	7(1)	14(1)	4.50
5. Na. acid prop. 14** oz/sack	3.5	1.8	0(0)	0(0)	0(0)	2(0)	9(0)	5.25
6. Na. acid prop. 7** oz/sack	3.5	1.0	0(0)	2(2)	9(6)	19(9)	22(15)	3.75
7. Na. acid prop. 12** oz/sack	3.5	1.4	0(0)	0(0)	16(7)	19(13)	22(14)	4.25
8. Na. acid prop. 7** oz/sack	3.4		4(2)	5(2)	16(3)	22(7)	22(18)	3.25
9. Na. acid prop. 12** oz/sack	3.5		0(0)	0(2)	6(8)	19(10)	19(15)	4.00 (3.50)

TABLE 4

d. Ca. prop. 9 oz/sack (280lb)	3.4	1.2	0(0)*	2(0)	8(2)	16(3)	20(5)	3.75
e. Ca. prop. 14 oz/sack "	3.5	1.5	0(0)	0(0)	0(0)	5(0)	12(1)	5.00
f. Ca. prop. 9 oz/sack "	3.5	1.0	0(0)	3(5)	7(9)	20(16)	21(19)	3.75
g. Ca. prop. 14 oz/sack "	3.5	1.7	0(0)	0(0)	8(4)	15(11)	18(15)	4.50
h. Ca. prop. 9 oz/sack "	3.5		2(2)	4(2)	11(6)	20(11)	22(14)	3.50
i. Ca. prop. 14 oz/sack "	3.6		0(0)	0(0)	6(0)	12(1)	16(3)	4.25

* Figures in brackets show cut surfaces mouldy.

** Weight of 70% aqueous solution per sack weighing 280 lbs.

5 The specific volume of the breads did not vary greatly in any of the tests, indicating that the additional yeast in the recipes was successful in maintaining volume. It is also clear from the data shown in Tables 1 and 2 that the use of acid propionates results in an improved oven spring in comparison with the full neutral salts.

10 The organoleptic panel was unable to detect any significant difference in odour between calcium propionate and 70% sodium dipropionate when used at similar levels in bread. It would appear that these two materials have a very similar odour threshold.

15 The results from the storage tests suggest that at similar concentrations the 70% sodium dipropionate formulation is more effective than solid calcium propionate in extending mould-free shelf life.

WHAT WE CLAIM IS:—

20 1. A method of inhibiting the growth of harmful micro-organisms in foodstuffs susceptible thereto comprising incorporating therein an acid propionate salt as hereinbefore defined in an amount of between 0.01 and 1.0% by the dry weight of the said foodstuff.

25 2. A method of inhibiting rope formation and mould growth in bread comprising incorporating in the bread an acid propionate salt in an amount of between 0.01 and 0.4% by the dry weight of the flour.

3. A method according to claim 2 wherein the bread is an enzyme leavened or chemically aerated bread.

4. A method according to claim 1 wherein the foodstuff is selected from cakes, pastry mixes, biscuits, flours, starches, sausages, meat pies and baby foods. 35

5. A method according to any of the preceding claims wherein the acid propionate salt is selected from sodium dipropionate, ammonium dipropionate, potassium dipropionate, calcium tetrapropionate and magnesium tetrapropionate. 40

6. A method according to any of the preceding claims wherein the acid propionate salt is added as an aqueous solution. 45

7. A process according to claim 2 wherein the acid propionate salt is added to the water used in the making of dough from flour in the bread making process. 50

8. A method of inhibiting the growth of harmful micro-organisms according to claim 1 or 2 as hereinbefore described with reference to the Examples.

9. Bread containing an acid propionate salt whenever produced by a process according to any of the preceding claims 2, 3 and 5—8. 55

J. HARRY,
Agent for the Applicants.

Printed for Her Majesty's Stationery Office by the Courier Press, Leamington Spa, 1978.
Published by the Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from which copies may be obtained.